

**FY 2015 Report on Monitoring and Management
of the Black-capped Vireo (*Vireo atricapilla*)
on Travis County's Balcones Canyonlands Preserve**



Photo: Black-capped Vireo hatchling (Paul Fushille)

**Travis County
Department of Transportation and Natural Resources
Natural Resources & Environmental Quality Division**



October 1, 2014 – September 30, 2015

TABLE OF CONTENTS

INTRODUCTION.....	4
STUDY SITES.....	6
HABITAT RESTORATION	8
METHODS	9
Data Collection	9
Banding	11
Data Analysis.....	11
RESULTS.....	12
DISCUSSION.....	178
RECOMMENDATIONS.....	22
LITERATURE CITED.....	25

LIST OF FIGURES

Figure 1. Balcones Canyonlands Preserve Location Map, Travis County, TX.....	29
Figure 2. BCVI Survey Locations and Observations 2015, Travis County, Texas.....	30
Figure 3. Restored BCVI Habitat and Habitat Restoration Areas (HMA) Travis, County, Texas (2015).....	31
Figure 4. BCVI Observations and Territories on the Jollyville Unit, Travis County, Texas (2015).....	32
Figure 5. BCVI Observations and Territories on the Ribelin and Sam Hamilton East tracts, Travis County, Texas (2015).....	33
Figure 6. BCVI Observations and Territories on the Kotrla Unit, Travis County, Texas (2015).....	34

LIST OF TABLES

Table 1. Summary of Jollyville Unit and Travis County BCVI survey data collected by various agencies, Travis County, Texas. 1989-2014.....	13
Table 2. Abundance, territory number, and pairing success rate for BCVIs (<i>Vireo atricapilla</i>) on Jollyville Unit, Travis County, Texas, March-September 2001-2015.....	14
Table 3. Breeding success and productivity for BCVI (<i>Vireo atricapilla</i>) territories on Jollyville Unit, Travis County, Texas, March-September 2001-2015.....	14
Table 4. Features of BCVI nests located on Travis County BCP lands in 2015.....	15
Table 5. BCVI (<i>Vireo atricapilla</i>) nest locations on Jollyville Unit, Travis County, Texas, March-September 2001-2015.....	17
Table 6. Proportion of second-year males (pSY) and site fidelity on Travis County BCP, Travis County, Texas. 2010-2015.....	18

INTRODUCTION

In the United States and Canada, 448 native bird species breed in terrestrial habitats (Rich et al. 2004). Approximately 200 of those terrestrial species, commonly known as neotropical migrants, breed in North America, and then migrate south to winter in Mexico, Central America, South America, and the Caribbean (Sibley 2001). A majority of neotropical migratory bird species face population declines due to a wide array of threats including, but not limited to, habitat loss, degradation, and fragmentation (Rich et al. 2004). Partners in Flight, the U.S. Fish and Wildlife Service (USFWS), and many other cooperating agencies are actively developing recovery and conservation plans, acquiring and protecting critical habitat, and educating the general public about bird conservation issues in order to slow or prevent further population declines.

The Black-capped Vireo (*Vireo atricapilla*, hereafter BCVI, or vireo) was listed as a federally endangered species in 1987. Habitat loss and nest parasitism by Brown-headed Cowbirds (*Molothrus ater*) are among the greatest threats to BCVI populations. The breeding range of this neotropical migrant has decreased markedly within the last few decades. Historically, BCVIs in the United States were found in Kansas, Oklahoma, and Texas during the breeding season (Grzybowski 1995). Presently, the BCVI breeds in a restricted range that includes: three counties in Oklahoma, portions of central and south-central Texas, and south into central Coahuila, through Nuevo Leon and into southwestern Tamaulipas, Mexico (Farquhar and Gonzalez 2005; Grzybowski et al. 1994; Grzybowski 1995, Oklahoma Dept. of Wildlife Cons. 1999; U.S. Fish and Wildlife Service 1991). Between 1996 and 2005, BCVI populations in Texas have been reported in only 38 south-central and central counties (Wilkins et al. 2006). A USFWS review of the BCVI population status was completed as required by the Endangered Species Act in 2006 (Wilkins et al. 2006).

Available BCVI habitat, and subsequently BCVI populations, in Travis County have been significantly reduced as natural disturbances, such as fire, are suppressed and suburban development continues to expand. Additionally, browse pressure from White-tailed deer (*Odocoileus virginianus*) overpopulation can negatively affect BCVI habitat structure (Grzybowski 1995). In 1996, less than 100 individual BCVIs were estimated to occur in Travis County (USFWS 1996a). More recent analyses of survey data indicate that since 2000, the BCVI population estimate in Travis County is fewer than 50 individuals (Wilkins et al. 2006).

The USFWS issued the Balcones Canyonlands Conservation Plan (BCCP) section 10(a)1(B) permit to the City of Austin and Travis County in 1996 (USFWS 1996b). The plan calls for a minimum of 30,428 acres of endangered species habitat in western Travis County to be set aside and managed within the Balcones Canyonlands Preserve (BCP or Preserve; see Figure 1). As of September 2015, the BCP encompassed over 31,700 acres. A minimum of 2,000 of these acres are required to be designated as BCVI habitat. Due to changes in land use and fire suppression, most BCVI habitat will have to be maintained and/or created through habitat restoration efforts (USFWS 1996c). Currently, Travis County manages over 8,800 acres designated part of the BCP.

BCVI habitat has been described as:

“low scrubby growth, mostly deciduous and of irregular height and distribution, with small spaces between the thickets and clumps, with vegetation cover to ground level. This is an early to mid-successional progression, or one maintained in edaphic (due to soil or topography) settings such as occurs in rocky gullies, edges of ravines, and on eroded slopes; thus often quite localized (Grzybowski 1995).”

Additionally BCVI habitat has been characterized as including the following:

- greater density of deciduous vegetation in height zones from 0 - 2 meters.
- average amounts of deciduous cover ranging from 30-45%, with total woody cover including Ashe juniper (*Juniperus Ashei*) ranging from 36-55%.
- greater within-territory heterogeneity of vegetation structure (with shrubs closely spaced but still separated, and allowing light to penetrate to ground levels).
- openness not exceeding about 65% in older adult vireo territories (i.e. at least 35% woody cover) (Grzybowski 1995, Bailey and Thompson 2007).

This report presents the results of the 2015 BCP surveys for BCVI conducted by Travis County Natural Resources personnel. Annual BCVI population data collected from Travis County-managed lands is analyzed in order to monitor changes in distribution, abundance, and productivity. Monitoring the population will help gauge the effectiveness of habitat restoration projects and prioritize sites for future restoration activities.

STUDY SITES

During the 2015 breeding season, BCVI territory surveys were primarily conducted on the Jollyville Unit (Cypress Creek Macrosite, BCP) and on the Ribelin tract (Bull Creek Macrosite, BCP) (Figure 2). Territory surveys were also conducted on the newly acquired Kotrla Unit (Big Sandy Creek Macrosite) in far northern Travis County (Figure 2). Additional secondary locations were checked periodically for presence/absence of BCVIs on other areas of the Jollyville Unit, the Ribelin tract and the Lake Travis Bluffs subsection of the Lucas tract (Figure 2).

The Jollyville Unit is comprised of 1,887 acres (764 ha) and includes the following tracts: Bunten, Collins, Cuevas, Cuevas East, Grandview Hills, Nootsie, Snowden, Vireo Ridge, and Vista Point (Figure 2). The Jollyville Unit is located approximately 13 miles (21 km) northwest of downtown Austin. Both Golden-cheeked Warbler (*Setophaga chrysoparia*, hereafter GCWA) and BCVI inhabit this unit. Topography includes upland plateaus that give way to irregular, steep slopes and ravines. Ravines drain into an unnamed tributary of Lake Travis (Colorado River) or into Cypress Creek. Drainages tend to flow from the east to west. The Travis County soil survey shows that riparian soils in these drainages are composed primarily of soils of the Volente complex; Brackett and Tarrant soils are found on steep slopes (USDA 1974). Tarrant soils also occur in level upland areas.

The Jollyville Unit contains closed canopy, oak-juniper (*Quercus sp.-Juniperus ashei*) woodlands, which cover the majority of the canyons and slopes. Historic harvest of mature Ashe juniper has allowed shrubby, secondary-growth junipers to dominate much the uplands and slopes. Open grasslands are found in some valleys and ridge tops, and riparian vegetation, which is dominated by black walnut (*Juglans nigra*), sycamore (*Platanus occidentalis*) and elms (*Ulmus spp.*), occupies riparian areas along creeks and drainages.

The Ribelin tract is located within the Bull Creek macrosite (Figure 2). This 319 acre (129.1 ha) tract is located off of RM 2222 and McNeil Drive, which is the primary access point. The tract is located between Travis County's Sam Hamilton East tract and the City of Austin's Kent Butler tract and the Upper Bull Creek Unit. The tract contains a short section of Bull Creek, and the topography includes upland plateaus that give way to irregular, steep slopes and ravines. Primary soils on this tract are found in the Tarrant series (USDA 1974). Closed canopy oak-juniper woodlands

cover the majority of the canyons and slopes. Humans have heavily impacted the lands comprising the Ribelin tract. There are several ranch roads, a substantial power line corridor (which makes up the south boundary of the property line), man-made clearings, old dumps, and fences found throughout the tract. In recent history, the land was utilized for cattle and livestock ranching.

The Kotrla Unit totals 950 acres (384 ha) is comprised of the Kotrla tract (500 acres) and the Scott & White tract (450 acres). These tracts are located in the Big Sandy Creek macrosite in far northwest Travis County. These properties contain habitat for both endangered songbirds and likely provide substantial connectivity between the BCP and BCNWR. The upland areas are relatively flat and are underlain by the hard gray limestone of the Edwards formation. These areas are classified as Karst Zone 2, and several known caves and sinks are scattered throughout the upland area.

Upland areas contain substantial Ashe juniper and areas of concentrated shin oak (*Quercus sinuata*) growing in shrub form. Other species include live oak (*Quercus virginiana* var. *fusiformis*), cedar elm (*Ulmus crassifolia*), Texas ash (*Fraxinus texensis*) and Texas oak (*Quercus fusiformis*). In particular, the north-eastern section of the property contains habitat currently suitable for the BCVI, as well as areas which might be made suitable by burning or other vegetative manipulation.

A substantial canyon runs generally northwest to southeast between the two tracts. The sloped areas are steep and heavily wooded. The tree canopy is generally taller than that on the upland areas, although the species composition is similar. There is a substantial deciduous component to the tree composition on these slopes, particularly Texas oak. A substantial colony of rare Texabama croton (*Croton alabamensis* var. *texensis*) occurs in this canyon.

The Lucas tract, which includes the Lake Travis Bluffs subsection, totals 297 acres (120 ha) located approximately 1.5 miles (2.4 km) northeast of Mansfield Dam which impounds Lake Travis (Colorado River). Primary access points are either on RM 620, which bounds the property to the south, or Comanche Trail, which bisects the property into eastern and western sections. The Lucas tract is part Lake Travis Unit of the Cypress Creek Macrosite.

Topography includes upland plateaus, steep slopes and ravines. Ravines drain directly into Lake Travis on the western portions of the property and into Bullick Hollow Creek, a tributary of Lake Travis, on the eastern portion. The Travis County

soil survey defines the surface soil types as part of the Brackett Association (USDA 1974). The uplands are dotted with karst features, including caves and sinks.

Vegetation types found on the Lucas tract are generally similar to those on the Jollyville Unit. Prior to Travis County ownership, small portions of this tract were cleared for livestock pens and hunting lanes. Malta starthistle (*Centaurea melitensis*), chinaberry (*Melia azedarach*) and Roosevelt weed (*Baccharis neglecta*) are common in these disturbed areas. The Lake Travis Bluffs section, acquired in 2011, is an 18.5 acre (7.5 ha) parcel that was previously cleared for development. Subsequently, most of the entire area has grown into excellent BCVI habitat dominated by shin oak sumacs (*Rhus* spp.) and Texas redbuds (*Cercis canadensis*).

HABITAT RESTORATION

Beginning in FY2001, and continuing every winter thereafter, Travis County has conducted BCVI habitat restoration within BCVI Habitat Management Areas (HMAs) (Figure 3). These HMAs are located within Travis County BCP lands and they have been designated as having the potential to be restored to BCVI habitat. HMA locations are targeted based on known historic vireo occupancy and soil and vegetative components necessary for vireo habitat. More HMAs will be developed as additional land with restoration potential is acquired.

Poor quality BCVI habitat has been altered through a variety of techniques to create the patchy, early to mid-successional physiognomy associated with BCVI breeding habitat. By selectively removing undesirable monoculture woody species such as Ashe juniper, the growth and shrubby structure of other woody species can be improved. The following woody species benefit from Ashe juniper removal: shin oaks, possumhaw (*Ilex decidua*), yaupon (*Ilex vomitoria*), Texas redbud (*Cercis Canadensis* var. *texensis*), wafer-ash (*Ptelea trifoliata*), Carolina buckthorn (*Frangula caroliniana*) and sumacs. Several sections of habitat, particularly those restored from 2001 to 2004, have become good to excellent quality habitat. It has been observed on the Jollyville HMA that BCVIs begin occupying restored areas three to five years following initial habitat manipulation.

During February 2015, 3.1 acres on the northern Vireo Ridge tract was targeted for restoration. This area expands upon a section originally restored in 2001 and where dense stands of shin oak had been choked out by Ashe juniper. All thinning of Ashe juniper and topping-off of shin oak was accomplished by hand. These actions

increased the cumulative total of restored habitat on Travis County managed lands to approximately 154.85 acres (62.67 ha).

Follow up clearing is required to maintain BCVI habitat at an early to mid-successional stage. In most previously manipulated areas, much of the refined hand clearing is completed in subsequent years rather than at the onset of initial larger scale brush removal which is typically completed mechanically. Portions of areas cleared in 2001-2003 are beginning to grow out of ideal BCVI habitat and will require additional maintenance in the near future. Other locations being considered for targeted habitat restoration in the future include areas on the Vireo Ridge, Lucas, Ribelin, New Life and Cuevas East tracts due to their history of occupancy and proximity to currently occupied habitat.

Restoration activities will continue where habitat potential has been identified on the HMAs. Target areas for restoration vary from year to year based primarily on the following: habitat utilized by BCVIs during the preceding breeding season, presence of occupied Golden-cheeked Warbler habitat, diversity of desired woody species, and available budget. Travis County staff utilizes a flexible multi-year BCVI habitat restoration plan which defines areas targeted for restoration and allows for additional areas to be added or removed based on new land acquisitions, changes in land use and updated management techniques and recommendations.

METHODS

Data Collection

Detailed protocol for BCVI territory mapping, nest monitoring, and presence/absence surveys are described in an unpublished report by Travis County (2009). In FY 2015, territory mapping was used to estimate BCVI abundance and number of individual territories present. All observations (both visual and auditory) of male, female and juvenile BCVIs were plotted on hard-copy, digital ortho-photo maps with a scale of 1:3,000 or less. The following data were recorded in the field for each observation: location, date, behavior, sex, age, presence of a mate, number of fledglings and color band combination (if banded). BCVI locations and corresponding data were later recorded into an ArcGIS 10 (ESRI, Inc., Redlands, CA) geodatabase with the Texas State Plane (with NAD 1983 projection) coordinate system. All males encountered were assigned a unique number (“territory number”) to signify them as

individuals. Female locations were either recorded as independent locations or with the corresponding territory number of a territorial male where applicable.

Territory mapping methods generally followed International Bird Census Committee (IBCC) guidelines (1970). Bibby's (2000) "consecutive flush" method was employed to increase accuracy in assigning observations of BCVIs to specific territories ("clusters"). However, Bibby's method was modified to be less intrusive in order to avoid harassment and/or human induced behaviors (Holiman and Craft 2000). No more than 10 locations for an individual bird were mapped at one time. The presence of returning color-banded BCVI allowed identification of particular individuals in many cases. When band status was unknown or when dealing with unmarked BCVI, conspecific singing or counter-singing was used to differentiate between males. BCVIs that could not be positively identified were designated as "unknown." Playback tapes of BCVI vocalizations were used infrequently late in the season in accordance with USFWS protocol to elicit BCVI responses.

Pairing status of male BCVIs was determined by observing one or more of the following conditions: a male associating with a female, an active nest associated with a male, and/or a male tending at least one fledgling. If any of the criteria for pairing success was met or if a male was observed in the same general location on at least three different occasions with at least a week between observations, it was considered to have established a breeding territory.

Nests were located opportunistically during normal territory mapping surveys. Disturbance was minimized by refraining from intensive nest searches. When found, nests were checked every five to seven days from the date of discovery until an outcome could be determined. Nest stage, contents, location and behavior of the attending adult were recorded during each nest check. In territories where no nest was found, but fledglings were present, staff recorded the maximum number of fledglings observed being attended. Fledglings observed being tended by a male or female confirmed breeding success within a territory; the total number of fledglings observed at any one time was used as a conservative measure of productivity.

In addition to territory mapping BCVIs on the Jollyville Unit, presence/absence surveys were conducted at the following seven additional sites (figure 2): 1) the east-west running ridge of the Vista Point tract (Jollyville Unit), referred to as "Coffee Cup Ridge" that was restored in 2008; 2) an area on the Ribelin tract that was occupied by a BCVI in 2012; 3) areas along a power line easement on the Ribelin tract that

were restored in 2009 and 2010 and occupied in 2013; 4) the power line easement on the Sam Hamilton tract; 5) a previously occupied area (2009 and 2010) on the Cuevas East tract; and 7) the Lake Travis Bluffs section of the Lucas tract (Lake Travis Unit) that was occupied in 2011. Survey effort on the Ribelin tract area actually exceeded what is required by the protocol as a result of the site being located on a GCWA survey plot. Subsequently, this location was visited more than the minimum required five visits directed by the survey protocol.

BCVIs were surveyed for a total of 179 hours from March 31 (first detection) to August 26, 2015 (last detection). Data can be interpreted as a comprehensive census due to the relatively large amount of time spent monitoring these birds.

Banding

In an effort to build a long term demographic data set, the banding program initiated in 2008 as part of a graduate study of avian dispersal, has continued. The original project investigated interpatch dispersal patterns within a fragmented preserve network (Simper 2009). Throughout the survey season, Travis County staff updates and shares data, including banded bird resightings and nest locations in an effort to coordinate banding attempts. In 2015, one adult male BCVI and one adult female BCVI were banded. Since 2008, 41 adult BCVIs have been banded on Travis County BCP properties. Travis County intends to continue the BCVI banding program initiated by this project.

Data Analysis

BCVI abundance is defined as the sum of all individual male BCVIs detected at a given study site (regardless of territory status). Overall species distribution is comprised of all locations where BCVIs were detected (i.e. registrations) and includes males, females, fledglings, and multiple sightings of the same individual. An individual male was considered to have established a breeding territory if the following behaviors were observed: 1) accompanying a female, 2) tending a nest or fledglings, or 3) singing in the same locality on three separate occasions each separated by one week (i.e. six days between observations). In calculating territory numbers, all BCVI territories, whether they were observed entirely or partially on Travis County property, were considered 'full' territories (as opposed to 'edge' or partial territories). Due to limitations of time and survey effort, the set of registrations shown on attached maps do not represent a definitive description of territory boundaries.

Pairing success rate was calculated as the proportion of territories within which a female was observed or a nest was located (Anders 2000). Productivity data is represented in the following two ways: 1) the total number of fledglings divided by the total number of territories and 2) the total number of fledglings divided by the number of successful territories. A territory was considered successful if at least one fledgling was observed with a territorial male or female. The breeding success rate is the proportion of full territories that successfully fledged young (Kolozsar and Becker 2000).

RESULTS

In 2015, BCVIs were located on the Jollyville Unit (Figure 4), the Ribelin and the Sam Hamilton East tracts of the Upper Bull Creek Unit (Figure 5) and the Kotrla Unit (Figure 6). Abundance totaled 21 males. Sixteen males established territories with 15 located in areas where habitat restoration had previously occurred. Five unique males did not establish territories, possibly being transient or migrating individuals, and were only accounted for in abundance. No BCVI territories were established in entirely new areas on the Jollyville Unit and none were detected in areas slotted for presence/absence surveys. Table 1 presents a summary of BCVI data collected during the 2015 breeding season in addition to data collected by various agencies exclusively on the Jollyville Unit since 1989. This table also indicates total abundance and territories on all Travis County properties (including BCVI detected from Travis County properties).

In 2015, eleven territories were considered successfully paired (Table 2). Seven territories fledged offspring, yielding at least 30 'hatch year' (HY) individuals observed in the field. Four territories each produced two successful nests. This was the first time since the monitoring program began that a territory definitively had more than one successful nest. Current protocol counts either the number of fledglings directly observed in the field or the number of nestlings discovered in a nest that is later demonstrated to be successful. Because of their dull-colored plumage, cryptic behavior, and unpredictable flights, we expect observer counts of fledglings to be biased low. Productivity in 2015 (Table 3), whether measured relative to successful territories (2.7 HY per terr.) or all territories (2.3 HY per terr.), is likely underestimated due to the fact that only six of eight (75%) territories with breeding success had nests that were monitored and thus exact HY totals were unknown.

Table 1. Summary of Jollyville Unit and Travis County BCVI survey data collected by various agencies, Travis County, Texas. 1989-2015.

Year Surveyed	Agency**	Jollyville Unit Abundance	Jollyville Unit No. Territories	Abundance/No. Territories (all Travis County lands)
1989	DLS	11	5	n/a
1990	DLS	11	5	n/a
1991	DLS	14	9	n/a
1992	TXDOT	Unknown ¹	Unknown	n/a
1993	TXDOT	25	20	n/a
1994	TXDOT	27	27	n/a
1995	TXDOT	23	23	n/a
1996	SWCA	19-22	15	n/a
1997	No Data	Unknown ¹	Unknown	n/a
1998	No Data	Unknown	Unknown	n/a
1999	No Data	Unknown	Unknown	n/a
2000	SWCA	3 ²	Unknown	n/a
2001	Travis County TNR	19	13	19/13
2002	Travis County TNR	21	19	21/19
2003	Travis County TNR	21	20	22/20
2004	Travis County TNR	13	12	13/12
2005	Travis County TNR	12	11	12/11
2006	Travis County TNR	12	9	13/9
2007	Travis County TNR	14	13	18/15
2008	Travis County TNR	15	15	15/15
2009	Travis County TNR	13 ³	11	15/11
2010	Travis County TNR	13	11	13/11
2011	Travis County TNR	9	8	10/9
2012	Travis County TNR	9	8 ⁴	9/8
2013	Travis County TNR	13	9	15/11
2014	Travis County TNR	14	12	14/12
2015	Travis County TNR	15	13	21/16

*Study area is only referred to as “Jollyville Unit” since 2001. ** See Literature Cited for appropriate report citation.

¹ Restricted access on the tract.

² Detected while on brief site visit.

³ Does not include individual detected on Lake Perspectives tract (now part of the Lake Travis Unit).

⁴ One banded male established two separate, isolated territories.

Table 2. Abundance, territory number, and pairing success for BCVIs (*Vireo atricapilla*) on Jollyville Unit, Travis County, Texas, March-September 2001-2015.

Year	Total Hours Surveyed ^a	Abundance	No. Territories	Successfully Paired	Pair Success Rate (%)
2001	100	18 ^a	15 ^a	13 ^a	86.7 ^a
2002	179	21	19	18	94.7
2003	290 ^b	21	19 ^a	14	73.7 ^a
2004	127.5	13	12	9	75
2005	140	12	11	8	72.7
2006	146	11 ^a	9	8	88.9
2007	178	14	13	12	92.3
2008	199	15	15	14	93.3
2009	204	13 ^c	11	11	100
2010	181	13	11	11	100
2011 ^d	131	9	8	8	100
2012	141	9	8	8	100
2013	154	13	9	9	100
2014	190	14	12	11	91.7
2015	179	15	13	11	84.6

^a Adjusted values to reflect BCVI found only on the JV Unit. • Total hours for entire season on all properties, not only JV Unit

^b Total hours surveyed for 2003 could not be verified and likely overestimated.

^c The male located at Lake Perspectives (Lake Travis Unit) was not included in this dataset.

^d The territory located at Lucas-Lake Travis Bluffs (Lake Travis Unit) was not included in this dataset.

Table 3. Breeding success and productivity for BCVI (*Vireo atricapilla*) territories on Jollyville Unit, Travis County, Texas, March-September 2001-2015.

Year	No. of territories with breeding success ¹	Breeding success rate (%)	No. Fledglings	No. fledglings/ successful territory ²	No. fledglings/ total no. territories ³
2001	4	26.7*	12	3.0*	0.8*
2002	13*	68.4*	25*	1.4*	1.3*
2003	9	47.4*	16	1.8	0.8
2004	6	50.0	13	2.2	1.1
2005	1	9.1	1	1.0	0.1
2006	8	88.9	15	1.9	1.7
2007	8	61.5	24	3.0	1.8
2008	10	66.7	29	2.9	1.9
2009	7	63.6	12	1.7	1.1
2010	8	72.7	10	1.3	0.9
2011	4	50.0	11	2.8	1.4
2012	7	87.5	21	3.0	2.6
2013	6	66.7	19	3.2	2.1
2014	8	66.7	23	2.9	2.1
2015	7	53.8	30	2.7	2.3

¹ Represents the number of all territories that fledged at least one young.

² Represents the average no. of fledgling from territories with breeding success.

³ Represents the average no. of fledgling from all successfully paired territories.

* Adjusted values to reflect BCVI found only on the JV Unit.

A total of 15 BCVI nests were located in 2015. One of these was discovered post-use (likely successful). Detailed data regarding nest substrate, height, and orientation were collected post breeding season for each nest found (Table 4). Table 4 also lists the outcome for each nest identified and Table 5 lists nest location data.

Of the nests monitored (including the one found post-fledge) [n=15], ten (66.7%) successfully fledged at least one young. Four nests were depredated (26.7%) by unknown predators with one of these abandoned while hatching due to mite infestation. Three territories had second nest attempts observed and monitored, that were successful. A fourth territory also had at least two successful nests with neither nest being located in time to monitor. Some initial nest discoveries occurred later in the season suggesting that earlier nests were never located and were likely unsuccessful.

Table 4. Features of BCVI (*Vireo atricapilla*) nests located on Travis County BCP in 2015.

Substrate	Primary substrate height (m)	Additional Concealment	Nest height (cm)	Distance from stem (cm)	Distance from foliar edge (cm)	Orientation	Comments
Mexican buckeye (<i>Ungnadia speciosa</i>)	2.5	n/a	46	96	12	SW 234	Successful; (3 HY)
Shin oak (<i>Quercus sinuata</i>)	5.0	Ashe juniper (<i>Juniperus Ashei</i>)	84	105	8	NW 316	Successful; (3 HY)
Texas ash (<i>Fraxinus texensis</i>)	9.0	Escarpment black cherry (<i>Prunus serotina</i> var. <i>eximia</i>)	119	46	23	SW 220	Failed; (2 eggs on ground)
Texas oak (<i>Q. fusiformis</i>)	10.0	Live oak (<i>Q. fusiformis</i>) Flameleaf sumac (<i>Rhus lanceolata</i>)	78	31	27	WSW 248	Failed; (4 nestlings)
Live oak	13.0	Flameleaf sumac	162	50	3	WSW 248	Failed second attempt; (3 eggs; mite infestation)

Substrate	Primary substrate height (m)	Additional Concealment	Nest height (cm)	Distance from stem (cm)	Distance from foliar edge (cm)	Orientation	Comments
Shin oak	4.0	n/a	38	24	5	SSE 158	Successful found post-fledge; (at least 1 HY)
Shin oak	3.0	n/a	92	25	19	N 10	Successful; (4 HY)
Shin oak	4.5	n/a	91	91	16	NNE 22	Successful; (4 HY)
Texas persimmon (<i>Diospyros texana</i>)	4.0	Shin oak	76	48	39	S 180	Successful second nest; (4 HY)
Shin oak	2.0	n/a	67	72	17	N 360	Failed (4 eggs); female also possibly depredated
Evergreen sumac (<i>Rhus virens</i>)	4.5	n/a	79	174	27	E 80	Successful; (2 nestlings, 2 unhatched eggs)
Evergreen sumac	1.5	Ashe juniper	83	124	14	ENE 64	Successful second nest, (1 HY, 1 unhatched egg)
Shin oak	3.0	Texas oak Dewberry (<i>Rubus riograndis</i>)	17	15	8	E 76	Successful; (4 HY)
Shin oak	4.0	Ashe juniper	20	24	10	SE 157	Successful; (3 HY)
Texas oak	3.0	Flameleaf sumac Nettleleaf hackberry (<i>Celtis laevigata</i> var. <i>reticulata</i>)	69	91	15	NW 312	Failed; (2 eggs)

Definitions: Substrate: plant species in which the nest is located. Concealment: plant species or other substrates that is primarily responsible for concealing the nest. Orientation: compass direction of nest relative to its substrate.

Table 5. BCVI (*Vireo atricapilla*) nest locations on Jollyville Unit, Travis County, Texas, 2015.

Object-ID	Color Band Combination	Nesting attempt	UTM E	UTM N	LAT	LONG
BLUE	DB/BK:DB/SI	1	610083	3368305	30.44179916	-97.85350037
BLUE	DB/BK:DB/SI	2	610113	3368319	30.44199944	-97.85320282
BRU	NB/SI:BK/BL	1	610120	3367909	30.43829918	-97.85320282
MR.P	PI/SI:NB/GR	1	609690	3368097	30.44000053	-97.85769653
VR06	UB	1	610229	3367757	30.43689919	-97.85209656
VR06	UB	2	610186	3367758	30.43689919	-97.85250092
BEVO	OR/DB:OR/SI	1	609528	3368213	30.44109917	-97.85929871
VR09	UB	1	610674	3367183	30.43169975	-97.84750366
TBRD	DG/SI:WH/DB	1	609740	3368380	30.44260025	-97.85710144
TBRD	DG/SI:WH/DB	2	609695	3368364	30.44239998	-97.85759735
RI01	UB	1	612595	3363735	30.40040016	-97.82790375
VR01	UB	1	609834	3368889	30.44709969	-97.85610199
VR03	UB	1	609861	3368553	30.44409943	-97.85579681
VR02	UB	1	609879	3368725	30.44569969	-97.85559845
VR02	UB	2	609798	3368598	30.44449997	-97.85649872

In all, a total of nine individual adult BCVIs (seven males, two females) wearing color bands were observed on Travis County properties in 2015. Of these, one adult male and one adult female were banded during the 2015 field season. Four additional male BCVIs (banded in 2014) and two additional female BCVI (banded in 2009 and 2014) were also observed this year. One male, banded in 2013 was briefly observed and soon disappeared never establishing a territory. It is worth mentioning that one banded female is A8Y (older than eight years old) and produced three young.

Adult males showed a 71.4% return rate (five of seven banded males observed in 2014). Four of the five returning males occupied the same area they held in 2014 with one claiming a new area approximately one quarter mile from his 2014 territorial area. As previously mentioned one banded male returned to his same 2014 territorial area but soon vanished without establishing a territory in 2015. Two of four banded females observed in 2014 returned in 2015, hence a 50% return rate of females. Nestlings were not banded due to permit restrictions.

Prior to 2008 there was no banding program in place, precluding staff from determining return rates, recruitment and associated data. From 2003 to 2009 BCVI territory and abundance numbers slowly declined despite a period of growth in 2007

and 2008. Only one male was banded in 2015 and it was a ‘second year’ (SY) male, i.e. males in their first reproductive season, and it established a territory. Due to the low sample size (only one newly banded pair), field observations were used to determine SY males based on all BCVI territories monitored. Table 6 summarizes relevant demographic information since 2010.

- Proportion of SY males (pSY): The proportion of second year males has been used as an indicator of recruitment. A minimum pSY value (29%) has been suggested as a measure of a stable population (USFWS 1996a). In 2015, the overall pSY, based on marked birds, was 16.6% (one of six). This figure is likely low due to the small sample size (one newly banded male). At least six additional BCVIs were noted as likely SYs based on observations in the field. Based on observations of only territorial males, pSY was unofficially at least seven of 16 (43.8%).
- Site fidelity: In 2015, five of seven banded males observed (banded) in 2014 returned indicating a 71.4% return rate on all Travis County properties. On the Jollyville Unit specifically, the return rate was 57.1% (four of seven). Females showed a 50% return rate (two of four, from 2014).

Table 6. Proportion of second-year males (pSY) and site fidelity on Travis County BCP, Travis County, Texas. 2010-2015.

Year	pSY	Site Fidelity (males)	Site Fidelity (females)
2010	27.3	70.0	n/a
2011	20.0*	25.0	n/a
2012	28.6*	66.7	50.0
2013	57.1	16.7	100.0
2014	62.5*	42.9	50.0
2015	43.8**	71.4	50.0

* conservative number, at least three (2011), one (2012) and two (2014) males were unsuccessfully aged.

** unofficial number based primarily on observations of territorial (n=7 of 16) males and not solely marked BCVIs.

DISCUSSION

The number of territories (+4) and overall abundance (+7) increased from the 2014 season’s totals, marking the largest overall increase (abundance and territory

number) since the 2006-2007 season. In the core vireo habitat on the Jollyville Unit, 12 males established territories in 2014. In 2015, there were 14 unique males detected with 13 establishing territories marking an increase of one territory. This marks the fifth year of stable or increasing numbers of territories and the third straight year of increasing territory numbers. Unlike 2014, several territories were discovered outside the Jollyville Unit. Two territories were again established on the Ribelin tract in 2015 as was the case in 2013, indicating that BCVI's often find suitable habitat underneath and along powerline corridors where vegetation is regularly kept in shorter conditions for maintenance and safety reasons. In the case of the Ribelin corridor, Travis County has itself restored one of these areas and worked with Austin Energy to restore a second area. The cause of the increase in BCVI numbers is unclear, but normal population fluctuations, the easing of the drought, changes in migration patterns through the area, and improved habitat could be factors.

On the Jollyville Unit, pairing success dropped below 90% for the first time since 2006, but was still high at nearly 85%. Overall productivity (2.7 fledged offspring per full territory) dropped slightly from the previous four years but remained one of the higher rates since 2001. Productivity for the total number of successfully paired territories (2.3 fledglings per full territory) increased from 2014 and was the second highest on record (since 2001).

For the first time on record for Travis County BCP, second successful nests for a territory were noted. Four individual territories produced successful second broods. Three of these were from territories where nests were located and monitored while the fourth territory's success was noted by either observing HYs in the field or upon discovering the just-fledged second nest. This territory was also the only territory where HYs were detected where no nest was monitored. These double broods helped offset those monitored nests that were depredated (n=5) and accounted for at least 20 HYs.

Although depredation of nests was generally low (26.7%), there was a strong indication that at least one female was depredated along with four young. Feathers were discovered directly under the nest and a female was never observed following this event for the territory. The male continued to defend the territory for two additional weeks before departing the area. Of note, the female that paired with this

male in 2014 was found paired in a proximate territory where she had two successful broods; albeit both nests were left with unfertilized eggs post-fledge.

Travis County acquired the tracts that comprise the Kotrla Unit for, among other things, its expansive swaths of potential BCVI habitat. Although most of the habitat has grown undesirable, small pockets remain and in one of these areas, a BCVI territory was monitored. A second unique male was also detected just off tract but not included in Figure 6, as it was never detected on Travis County property. In 2014, a BCVI was also detected in a third location during a site visit prior to the County's purchase. The nearby location to the USFWS Balcones National Wildlife Refuge and its's substantial BCVI population makes this new unit important to the regional stability of this species. Habitat restoration efforts will likely begin during the winter of 2016-17 after a season of baseline data is acquired during the 2016 breeding season.

Total survey effort (11.2 hours/territory) decreased from the 2014 effort (15.8 hours/territory) and was a decrease from the overall survey effort average (15.7 hours/territory) from the previous five seasons (2010-2014). The amount of survey hours in 2015 decreased from 2014 by 11 hours. This decrease in effort can be attributed to personnel shortage and the increase in abundance and territories that required monitoring. An increase in survey and banding effort in 2016 is both likely (with the addition of staff) and warranted. The cryptic behavior of certain individual males also made it difficult to determine territory status (pairing and productivity). Survey hours are more directly tied to the survey team's ability to acquire pairing and productivity data (nests) than on BCVI abundance and numbers of territories. It is important to note that survey hours also include those in areas without BCVIs (presence/absence surveys).

Several factors may influence territory distribution, including the intrinsic habitat characteristics of the site, the age structure of the population, overall population density, and habitat restoration activities in protected areas (Grzybowski et al. 1994; Anderson and Gutzwiller 1996). Territory establishment in 2015 occurred in the same general core areas on the Jollyville Unit as in previous years (Travis County 2001-2014). Although the same areas are being utilized, often by returning males (documented by resighting of banded birds), the distribution of BCVI territories on the Jollyville Unit has changed each year. An area of special note is a portion of Vireo Ridge referred to as "35-acres". This portion was occupied by two BCVIs in

2003 and 2004. In 2005 a lone BCVI was detected once and but did not establish a territory. Because this area was growing out of habitat it was restored in winter 2005. It appeared to have grown back into suitable habitat by 2009 and was finally re-occupied by a successfully paired (and banded) male in 2013. In 2014, two males established territories in this area and one male again established a territory in 2015. Similarly, areas on the southern edge of the Jollyville core area have increased to three territories in 2015 after several years with no territories.

Habitat loss and degradation is a primary concern for BCVI recruitment onto Travis County BCP properties. Typical BCVI nesting habitat was uncommon on the Jollyville Unit when first acquired by Travis County in 2000 and 2001. Habitat change through vegetational succession, dominated primarily by encroaching Ashe juniper, reduced canopy openings. Additionally, the successional change in vegetation structure tends towards closed canopy woodlands, rather than the low, shrubby, mid-successional stage preferred by BCVIs. Through mechanical efforts of BCVI habitat restoration, significant portions of the Jollyville Unit HMA now show characteristics of prime BCVI habitat.

All of the 2015 BCVI territories on the Jollyville Unit, as well as those on the Ribelin utility corridor, were established primarily in previously manipulated areas which may be considered “restored”. This has been the observed trend since 2006. These observations indicate that BCVIs continue to respond positively to habitat restoration efforts undertaken since 2001, when it appeared that the remaining BCVIs were shifting annually from lesser quality habitat into higher-quality restored areas. It should be noted, however, that large areas of apparently restored habitat still have not been reoccupied and other areas that, although have been occupied, appear to have enough space for additional territories. Numerous factors influence territory establishment, but regional natural population fluctuations, declining connectivity, as well as the location of the Preserve on the far eastern edge of BCVI range are all likely to contribute to overall population trends. It is worth mentioning that although the last two years have generally been wetter, the general prolonged drought over the last few years is likely negatively impacting the population. These impacts may include decreased habitat suitability e.g., decreased invertebrate population or sparse vegetation cover, which in turn could affect reproductive success.

In general, BCVIs tend to be “semi-colonial” and establish territories near other BCVIs (Ward and Schlossberg 2001). However, isolated territories composed of

either individual pairs or small-clusters have occasionally been located (Grzybowski 1990). These territories are often difficult to detect as noted by Ward and Schlossberg (2001). They found that in low-density populations of BCVIs, song rates and duration of song bouts were much lower than in high-density populations. Thus, low detection rates of these isolated territories may affect estimates of abundance. This can be noted on the Sam Hamilton East tract where two unique males were detected later in the season in a heavily wooded canyon. Although they may have been migrating, it is possible that they had established, or attempted to establish, territories on nearby ridges and were never detected due to their isolation.

Continued research and monitoring of BCVI on the Jollyville Unit is essential to determine whether the colony is a source population or a population sink. Source populations contribute to the general overall population and help to maintain satellite colonies. Population sinks are maintained solely through immigration and contribute nothing to species-wide abundance (Pulliam 1988). Data obtained from continued banding efforts would improve estimates of inter-population connectivity as well as age-specific survival and reproduction. If adequate connectivity with neighboring sub-populations is not maintained, then BCVI numbers on isolated preserve tracts may decline precipitously, leading to local extinctions and deficient levels of colonization. Information regarding productivity and dispersal is limited because it is difficult and slow to obtain. Additional survey seasons and continued research will help gauge the effectiveness of habitat restoration projects and prioritize sites for future restoration. Without comprehensive productivity, survivorship, and dispersal information, as well as complimentary data from surrounding properties, we cannot make a definitive statement regarding the role of the Jollyville Unit within the central Texas portion of the BCVI's range.

RECOMMENDATIONS

- Continue to conduct intensive BCVI monitoring, including annual color-banding. Such research could help address many questions concerning the long-term viability of the Jollyville Unit colony and its relationship to other sub-populations in the region. The long-term data sets generated from such studies would facilitate yearly comparisons and improve the accuracy of model-generated population forecasts. Data collected should include the following: abundance, age structure, dispersal patterns, distribution of subpopulations, habitat associations throughout the breeding season, nest success, recruitment, return rates and age-specific measures of productivity

and survivorship. Research should be focused upon answering questions of immediate relevance to management and recovery priorities.

- Collect and compile age-specific survivorship and reproductive information for marked birds. This information may provide a more accurate picture of population dynamics in cases where field data is believed to be incomplete or biased (e.g. using mean values to account for number of HYs per nest rather than number of fledglings observed in the field).
- Continue to facilitate opportunities for graduate students to collect and analyze quantitative habitat and vegetation data in order to construct habitat suitability and dispersal models. Longitudinal studies are needed to compare the results of habitat restoration treatments over time. Ideally, such data would be collected in a GIS-compatible format and at a scale that would allow preserve-wide spatial analysis. Such analyses could improve our ability to locate existing areas of high quality habitat as well as areas with high potential for successful restoration. Additionally, further research is needed to better understand dynamics of vegetative succession and its relationship to BCVI habitat suitability and reproductive success in this region.
- Continue to refine and standardize BCVI monitoring procedures, giving special attention to minimum hours of survey and nest searching needed. This is especially important for new properties with potential BCVI habitat.
- BCVI habitat and potential habitat for restoration should continue to be identified and mapped on all tracts owned and managed by Travis County. Historic BCVI locations on properties owned and managed by Travis County should be visited regularly during the field season to determine presence/absence of BCVIs.
- Habitat restoration efforts will continue on tracts that support BCVI populations and on surrounding tracts that harbor potential habitat. Restoration methods will be evaluated to determine the best techniques for creating suitable BCVI breeding habitat.
- Create a more structured GCWA survey methodology for monitoring territories and productivity in both potential and managed BCVI habitat. Efforts should be made to document any co-occurrences of GCWA and BCVIs during the breeding season in order to investigate and evaluate the possibility of creating areas of mixed or composite habitat.

- Judicious use of recorded BCVI vocalizations should continue in accordance with USFWS protocols, especially in areas where few and/or isolated BCVIs have been detected. This recommendation is also useful to determine absence of a target species. Playback should increase detections of territorial males (Horne 2000).
- Cowbird trapping should continue on and adjacent to any Travis County-managed properties occupied by endangered songbirds, with traps added or removed based on cowbird activity. Shooting female cowbirds in habitat is also recommended. The addition of the Kotrla Unit deep within rural northwestern Travis County will likely demand an aggressive trapping program in that area. Joining efforts with the USFWS Balcones National Wildlife Refuge or volunteers should be considered.
- Although it was unclear if any BCVI nest attempts failed due to red imported fire ant (*Solenopsis invicta*, RIFA) predation in 2015, control of this nuisance species should continue within restored BCVI habitat areas when warranted. RIFA have been documented as a common cause of nest failure in BCVI (Smith, et al. 2002). At two nest locations in 2015, fire ant mounds were treated within 25 m of the nest. Twenty-five meters is based on the foraging distance of RIFA (USFWS 2014).

LITERATURE CITED

- Anders, A. D. 2000. Demography of golden-cheeked warblers on Fort Hood, Texas in 2000: productivity, age structure, territory density, and adult return rates. In Endangered species monitoring and management at Fort Hood, Texas: 2000 annual report. Fort Hood Project, The Nature Conservancy of Texas, Fort Hood, Texas, USA.
- Anderson, S. H., and K. J. Gutzwiller. 1996. Habitat evaluation methods. Pages 592-606 in T. A. Bookhout, ed. Research and management techniques for wildlife and habitats. Fifth ed., rev. The Wildlife Society, Bethesda, Maryland, USA.
- Balcones Canyonlands Preserve Land Management Plan 2007. BCVI management. Chapter 8 in Tier II A, Balcones Canyonlands Management Handbook. Unpublished document. Travis County, Texas, USA.
- Bailey, W. J., and F. R. Thompson. 2007. Multiscale Nest-Site Selection by BCVIs. *Journal of Wildlife Management* 71:828-836.
- Bibby, C. J., N. D. Burges, D. A. Hill and S. H. Mustoe. 2000. Bird census techniques. Second edition. Academic Press.
- DLS Associates. 1989b. BCVI monitoring program, Travis County, Texas and vicinity, 1989. Submitted to: The Nature Conservancy of Texas, Inc. and Executive Committee of the Austin Regional Habitat Conservation Plan.
- _____. 1990a. BCVI monitoring program, Travis County, Texas and vicinity, 1990. Submitted to: The Nature Conservancy of Texas, Inc. and Executive Committee of the Balcones Canyonlands Habitat Conservation Plan.
- _____. 1992. BCVI monitoring program, Travis County, Texas and vicinity, 1991. Submitted to: The Nature Conservancy of Texas, Inc. and Executive Committee of the Balcones Canyonlands Habitat Conservation Plan.
- Farquhar, C.C. and J. I. Gonzalez. 2005. Breeding habitat, distribution and population status of the BCVI in northern Mexico. Project WER65 Final Report. U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- Grzybowski, J.A. 1990. Population and nesting ecology of the Black-capped vireo--1990. Austin (TX): Texas Parks and Wildlife Department. Report nr E-1-3. Job 29. 39 p.
- _____. 1995. BCVI (*Vireo atricapillus*). In *The Birds of North America*, No. 181 (A. Poole and F. Gill, editors). The Academy of Natural Sciences, Philadelphia, Pennsylvania, USA and The American Ornithologists' Union, Washington, D.C., USA.
- _____, D. J. Tazik, and G. D. Schnell. 1994. Regional analysis of BCVI breeding habitats. *Condor* 96: 512-544.
- Holiman, W. C., and R. A. Craft. 2000. Breeding density and productivity of the golden-cheeked warbler on Fort Hood, Texas in 1999. Pages 79-103 in *Endangered Species monitoring and management at Fort Hood, Texas: 1999 annual report*. Fort Hood Project, The Nature Conservancy of Texas, Fort Hood, Texas.

- Horne, J. S. 2000. Use of conspecific taped-playback to increase detectability of BCVI. Endangered species monitoring and management at Fort Hood, Texas: 1999 annual report. Revised edition. Fort Hood Project, The Nature Conservancy of Texas, Fort Hood, Texas, USA.
- International Bird Census Committee. 1970. An international standard for a mapping method in bird census work, recommended by the International Bird Census Committee. Audubon Field Notes 24 (6):722-726.
- Koloszar, J. A. and H. M. Becker. 2000. Monitoring of the BCVI during 2000 on Fort Hood, Texas. Endangered species monitoring and management at Fort Hood, Texas: 2000 annual report. Fort Hood Project, The Nature Conservancy of Texas, Fort Hood, Texas, USA.
- Oklahoma Department of Wildlife Conservation. 1999. Reproductive enhancement and population monitoring of BCVIs in Blaine County. Federal Aid Project E-45-2. May 1, 1998-April 30, 1999. Oklahoma Department of Wildlife Conservation. Oklahoma City, Oklahoma, USA. 14 p.
- Pulliam, H.R. 1988. Sources, sinks, and population regulation. American Naturalist 132: 652-661.
- Rich, T. D., C. J. Beardmore, H. Berlanga, P. J. Blancher, M. S. W. Bradstreet, G. S. Butcher, D. W. Demarest, E. H. Dunn, W. C. Hunter, E. E. Iñigo-Elias, J. A. Kennedy, A. M. Martell, A. O. Panjabi, D. N. Pashley, K. V. Rosenberg, C. M. Rustay, J. S. Wendt, T. C. Will. 2004. Partners in Flight North American Landbird Conservation Plan. Cornell Lab of Ornithology. Ithaca, NY. Partners in Flight website. http://www.partnersinflight.org/cont_plan/ (VERSION: March 2005).
- Sibley, D. A. 2001. The Sibley Guide to Bird Life and Behavior. Alfred A. Knopf, Inc. New York. 587 p.
- Simper, W.S. 2009. Black-capped vireo (*Vireo atricapilla*) color banding and surveying on the Balcones Canyonlands Preserve and the Balcones Canyonlands National Wildlife Refuge: Project Year 2. Austin (TX): USFWS. Report nr TE168189-0-2009. 32 p.
- Smith, J.E., S.J. Taylor and C.J. Whelan. 2002. Red Imported Fire Ant (*Solenopsis invicta*) depredation of Black-capped Vireo (*Vireo atricapillus*) nests at Ft. Hood, Texas. Illinois Natural History Survey Technical Report 2002(24).
- SWCA 1996. Master BCVI survey map, Grandview Hills Parcel 13. Unpublished document. Travis County, Texas, USA.
- Texas Department of Transportation. 1992. BCVI monitoring program, northwest Travis County, Texas. Unpublished report.
- _____. 1993. BCVI monitoring program, northwest Travis County, Texas. Unpublished report.
- _____. 1994. BCVI monitoring program, northwest Travis County, Texas. Unpublished report.

- _____. 1995. BCVI monitoring program, northwest Travis County, Texas. Unpublished report.
- _____. 2001. Monitoring of the Golden-cheeked warbler 2001 field season. Unpublished report.
- _____. 2001. Status of the BCVI on the Vireo Ridge Tract, Travis County, Texas, 2001 field season. Unpublished report.
- _____. 2002. Monitoring of the BCVI, 2002 field season, Travis County, Texas. Unpublished report.
- _____. 2003. Monitoring of the BCVI, 2003 field season, Travis County, Texas. Unpublished report.
- _____. 2004. Monitoring of the BCVI, 2004 field season, Travis County, Texas. Unpublished report.
- _____. 2005. Monitoring of the BCVI, 2005 field season, Travis County, Texas. Unpublished report.
- Travis County Natural Resources and Environmental Quality Division. 2006. Monitoring of the BCVI, 2006 field season, Travis County, Texas. Unpublished report.
- _____. 2007. Monitoring of the BCVI, 2007 field season, Travis County, Texas. Unpublished report.
- Travis County Natural Resources and Environmental Quality Division. 2008. Monitoring and Management of the BCVI (*Vireo atricapilla*) in Western Travis County, Texas: 2008 field season, Travis County, Texas. Unpublished report.
- _____. 2009. Monitoring and Management of the BCVI (*Vireo atricapilla*) in Western Travis County, Texas: 2009 field season, Travis County, Texas. Unpublished report.
- _____. 2010. Monitoring and Management of the BCVI (*Vireo atricapilla*) in Western Travis County, Texas: 2010 field season, Travis County, Texas. Unpublished report.
- _____. 2011. Monitoring and Management of the BCVI (*Vireo atricapilla*) in Western Travis County, Texas: 2011 field season, Travis County, Texas. Unpublished report.
- _____. 2012. Monitoring and Management of the BCVI (*Vireo atricapilla*) in Western Travis County, Texas: 2012 field season, Travis County, Texas. Unpublished report.
- _____. 2013. Monitoring and Management of the BCVI (*Vireo atricapilla*) in Western Travis County, Texas: 2013 field season, Travis County, Texas. Unpublished report.
- _____. 2014. Monitoring and Management of the BCVI (*Vireo atricapilla*) in Western Travis County, Texas: 2014 field season, Travis County, Texas. Unpublished report.
- U.S. Department of Agriculture. 1974. Soil survey of Travis County, Texas. United States Department of Agriculture, Soil Conservation Service, Washington, D.C. 123pp.
- U.S. Fish and Wildlife Service. 1991. BCVI population and habitat viability assessment report. Compiled and edited by Carol Beardmore, Jeff Hatfield, and Jim Lewis in conjunction with workshop participants. Report of a September 18-21, 1995

- workshop arranged by the USFWS in partial fulfillment of U.S. National Biological Service Grant No. 80333-1423. Austin, Texas. Pp ix + 57.
- _____. 1996a. BCVI (*Vireo atricapillus*) recovery plan. U.S. Fish and Wildlife Service, Austin, Texas, USA.
- _____. 1996b. Federal Fish and Wildlife Permit No. PRT-788841.
- _____. 1996c. Habitat Conservation Plan and Final Environmental Impact Statement. U.S. Fish and Wildlife Service, Albuquerque, New Mexico, USA.
- _____. 2014. Karst Preserve Managing and Monitoring Recommendations. Austin Ecological Services Field Office, Austin, TX. Pg. 14.
- Ward, M. and S. Schlossberg. 2001. Conspecific attraction in BCVIs. 2001 Annual Report. Natural Resources Branch. Endangered Species Management Program, Ft. Hood, Texas
- Wilkins, N., R. A. Powell, A. A. T. Conkey and A. G. Snelgrove. 2006. Population Status and Threat Analysis for the BCVI. Department of Wildlife and Fisheries Sciences, Texas A&M University.





Figure 1: Balcones Canyonlands Preserve Location Map Travis County, TX.



Legend

-  Jollyville Unit
-  Kotrla Unit
-  BCP
-  Roads
-  Rivers & Lakes
-  BCNWR

